


Sub
AT E1
--This application is a divisional of prior application
U.S. Serial No. 09/938,602, filed September 26, 1997, ^{now U.S. Pat. No. 6,380,194} which
claims priority from United States provisional application
Serial No. 60/027,260, filed October 1, 1996, the contents of
which are hereby incorporated by reference.--

In the Claims:

Cancel claims 5, 7-10, 13 and 15 without prejudice, amend
claims 1-4, 6, 11-12, 14 and 16-17, and add new claim 18 as
follows.

1. A compound of formula



a pharmaceutically acceptable acid addition salt or a
stereochemically isomeric form thereof, wherein

R¹ and R² are each independently selected from hydrogen;
hydroxy; amino; C₁₋₆alkyl; C₁₋₆alkyloxy; C₁₋₆alkylcarbonyl;
C₁₋₆alkyloxycarbonyl; Ar¹; mono- or di(C₁₋₆alkyl)amino;
mono- or di(C₁₋₆alkyl)aminocarbonyl; dihydro-2(3H)-
furanone; C₁₋₆alkyl substituted with one or two substituents
each independently selected from amino, imino,
aminocarbonyl, aminocarbonylamino, hydroxy, hydroxyC₁₋₆
alkyloxy, carboxyl, mono- or di(C₁₋₆alkyl)amino,
C₁₋₆alkyloxycarbonyl and thienyl; or

R¹ and R² taken together may form pyrrolidinyl, piperidinyl,
morpholinyl, azido or mono- or di(C₁₋₆alkyl)aminoC₁₋₆
alkylidene;

R³ is hydrogen, Ar¹, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆
alkyloxycarbonyl, C₁₋₆alkyl substituted with C₁₋₆
alkyloxycarbonyl; and

R⁴, R⁵, R⁷ and R⁸ are each independently selected from
hydrogen, hydroxy, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano,


aminocarbonyl, nitro, amino, trihalomethyl or trihalomethyloxy ;

R⁶ is aminocarbonyl; or

L is C₁₋₁₀alkyl substituted with one or two substituents independently selected from C₃₋₇cycloalkyl; indolyl or indolyl substituted with one, two, three or four substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C₁₋₆alkylcarbonyl; phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C₁₋₆alkylcarbonyl; and,

Ar¹ is phenyl; or phenyl substituted with one, two or three substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, nitro or trifluoromethyl.

2. A compound according to claim 1 wherein R¹ and R² are each independently selected from hydrogen, C₁₋₆alkyl, Ar¹ or mono- or di(C₁₋₆alkyl)aminocarbonyl; or R¹ and R² taken together may form pyrrolidinyl, piperidinyl or morpholinyl; R³ is hydrogen, C₁₋₆alkyl or Ar¹; and Ar¹ is phenyl, or phenyl substituted with one, two or three substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, nitro or trifluoromethyl; and L is a radical of formula



wherein Alk is C₁₋₆alkanediyl;

R^a, R^b, R^c, R^d, R^e, R⁴, R⁵, R⁷ and R⁸ are each independently selected from hydrogen, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl or trihalomethyloxy; or

R^a and R^b taken together may form a bivalent radical of formula

710340

-CH=CH-NR⁹- (a-1),

-NR⁹-CH=CH- (a-2),

wherein R⁹ is hydrogen or C₁₋₄alkyl.

3. A compound according to claim 2 wherein L is C₃₋₁₀alkenyl or C₁₋₂alkyl substituted with one or two substituents independently selected from C₃₋₇cycloalkyl; indolyl or indolyl substituted with one, two, three or four substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C₁₋₆alkylcarbonyl; phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C₁₋₆alkylcarbonyl.


4. A compound according to claim 3 wherein L is 2,6-dichlorophenylmethyl.

5. A compound according to claim 4 wherein NR¹R² is other than amino.

6. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically active amount of a compound as claimed claim 1.

12. A process for preparing a pharmaceutical composition comprising intimately mixing a therapeutically effective amount of a compound as claimed in claim 1 with a pharmaceutically acceptable carrier.

14. The combination of a compound of formula (I)



wherein R^1 and R^2 are each independently selected from hydrogen; hydroxy; amino; C_{1-6} alkyl; C_{1-6} alkyloxy; C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl; Ar^1 ; mono- or di(C_{1-6} alkyl)amino; mono- or di(C_{1-6} alkyl)aminocarbonyl; dihydro-2(3H)-furanone; C_{1-6} alkyl substituted with one or two substituents each independently selected from amino, imino, aminocarbonyl, aminocarbonylamino, hydroxy, hydroxy C_{1-6} alkyloxy, carboxyl, mono- or di(C_{1-6} alkyl)amino, C_{1-6} alkyloxycarbonyl and thienyl; or R^1 and R^2 taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C_{1-6} alkyl)amino C_{1-6} alkylidene;

R^3 is hydrogen, Ar^1 , C_{1-6} alkylcarbonyl, C_{1-6} alkyl, C_{1-6} alkyloxycarbonyl, C_{1-6} alkyl substituted with C_{1-6} alkyloxycarbonyl; and


R^4 , R^5 , R^7 and R^8 are each independently selected from hydrogen, hydroxy, halo, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl or trihalomethoxy;

R^6 is selected from cyano or aminocarbonyl;

L is C_{1-10} alkyl; C_{3-10} alkenyl; C_{3-10} alkynyl; C_{3-7} cycloalkyl; or L is C_{1-10} alkyl substituted with one or two substituents independently selected from C_{3-7} cycloalkyl; indolyl or indolyl substituted with one, two, three or four substituents each independently selected from halo, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethoxy, C_{1-6} alkylcarbonyl; phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethoxy, C_{1-6} alkylcarbonyl; and,

Ar^1 is phenyl, or phenyl substituted with one, two or three substituents each independently selected from halo, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, nitro or trifluoromethyl; and another antiretroviral compound.

16. A product containing (a) a compound of formula (I)



wherein R¹ and R² are each independently selected from hydrogen; hydroxy; amino; C₁₋₆alkyl; C₁₋₆alkyloxy; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; Ar¹; mono- or di(C₁₋₆alkyl)amino; mono- or di(C₁₋₆alkyl)aminocarbonyl; dihydro-2(3H)-furanone; C₁₋₆alkyl substituted with one or two substituents each independently selected from amino, imino, aminocarbonyl, aminocarbonylamino, hydroxy, hydroxC₁₋₆alkyloxy, carboxyl, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyloxycarbonyl and thienyl; or

R¹ and R² taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C₁₋₆alkyl)aminoC₁₋₆alkylidene;

R³ is hydrogen, Ar¹, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkyl substituted with C₁₋₆alkyloxycarbonyl; and

R⁴, R⁵, R⁷ and R⁸ are each independently selected from hydrogen, hydroxy, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl or trihalomethyloxy;


R⁶ is selected from cyano or aminocarbonyl;

L is C₁₋₁₀alkyl; C₃₋₁₀alkenyl; C₃₋₁₀alkynyl; C₃₋₇cycloalkyl; or

L is C₁₋₁₀alkyl substituted with one or two substituents independently selected from C₃₋₇cycloalkyl; indolyl or indolyl substituted with one, two, three or four substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C₁₋₆alkylcarbonyl; phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C₁₋₆alkylcarbonyl; and,

Ar¹ is phenyl, or phenyl substituted with one, two or three substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, nitro or trifluoromethyl; and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in anti-HIV treatment.

17. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and as active ingredients (a) a compound of formula (I)



wherein R¹ and R² are each independently selected from hydrogen; hydroxy; amino; C₁₋₆alkyl; C₁₋₆alkyloxy; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; Ar¹; mono- or di(C₁₋₆alkyl)amino; mono- or di(C₁₋₆alkyl)aminocarbonyl; dihydro-2(3H)-furanone; C₁₋₆alkyl substituted with one or two substituents each independently selected from amino, imino, aminocarbonyl, aminocarbonylamino, hydroxy, hydroxC₁₋₆alkyloxy, carboxyl, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyloxycarbonyl and thienyl; or R¹ and R² taken together may form pyrrolidinyl, piperidinyl, morpholinyl, azido or mono- or di(C₁₋₆alkyl)aminoC₁₋₄alkylidene; R³ is hydrogen, Ar¹, C₁₋₆alkylcarbonyl, C₁₋₆alkyl, C₁₋₆alkyloxycarbonyl, C₁₋₆alkyl substituted with C₁₋₆alkyloxycarbonyl; and R⁴, R⁵, R⁷ and R⁸ are each independently selected from hydrogen, hydroxy, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl or trihalomethoxy; R⁶ is selected from cyano or aminocarbonyl; L is C₁₋₁₀alkyl; C₃₋₁₀alkenyl; C₃₋₁₀alkynyl; C₃₋₇cycloalkyl; or

AL
L is C₁₋₁₀alkyl substituted with one or two substituents independently selected from C₃₋₇cycloalkyl; indolyl or indolyl substituted with one, two, three or four substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C₁₋₆alkylcarbonyl; phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, aminocarbonyl, nitro, amino, trihalomethyl, trihalomethyloxy, C₁₋₆alkylcarbonyl; and, Ar¹ is phenyl, or phenyl substituted with one, two or three substituents each independently selected from halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, nitro or trifluoromethyl; and (b) another antiretroviral compound.

18. A method of treating a subject suffering from HIV (Human Immunodeficiency Virus) infection comprising administering to the subject a therapeutically effective amount of the compound of claim 1.